

## University of Groningen

### Comment on Salawu et al.

Meulenbelt, H.E.; Geertzen, J.H.B.

*Published in:*  
Prosthetics and Orthotics International

*DOI:*  
[10.1080/0309364071300928](https://doi.org/10.1080/0309364071300928)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2007

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Meulenbelt, H. E., & Geertzen, J. H. B. (2007). Comment on Salawu et al. *Prosthetics and Orthotics International*, 31(2), 207-208. <https://doi.org/10.1080/0309364071300928>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).


The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

This article was downloaded by:[University of Groningen]  
On: 24 September 2007  
Access Details: [subscription number 770299803]  
Publisher: Informa Healthcare  
Informa Ltd Registered in England and Wales Registered Number: 1072954  
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Prosthetics and Orthotics International

Publication details, including instructions for authors and subscription information:  
<http://www.informaworld.com/smpp/title-content=t714595820>

### Comment on Salawu et al.

Henk E. Meulenbelt <sup>a</sup>; Jan H. B. Geertzen <sup>a</sup>

<sup>a</sup> Center for Rehabilitation, University Medical Center Groningen and Northern  
Center for HealthCare Research, University of Groningen, The Netherlands

Online Publication Date: 01 June 2007

To cite this Article: Meulenbelt, Henk E. and Geertzen, Jan H. B. (2007) 'Comment on Salawu et al.', Prosthetics and Orthotics International, 31:2, 207 - 208

To link to this article: DOI: 10.1080/03093640701300928

URL: <http://dx.doi.org/10.1080/03093640701300928>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## LETTER TO THE EDITOR

### Comment on Salawu et al.

Sir: In the December 2006 issue of *Prosthetics and Orthotics International* Salawu et al. provide us with very important information regarding stump ulcers. Stump ulcers are a frequently seen problem in amputees wearing prosthesis. These ulcers can disrupt daily activities and can therefore interfere with activities and participation of amputees. A study into whether prosthesis can be worn despite having ulcers is indeed important. The authors came to a very important conclusion, namely that allowing amputees with stump ulcers to continue using their prostheses is safe. This is, however, also a study which may lead to controversy, because no evidence is available in medical literature which supports one of the mentioned solutions namely leaving the prosthesis off until proper healing has occurred compared with continuing use of the prosthesis (with adding liners or socks).

We have some remarks on this conclusion.

As the authors state, their measure of prosthetic limb usage was (too?) crude and the registration of appropriate socket modifications was insufficient. It is our opinion that correction of this would be the first appropriate step in treating stump ulcers whether or not in combination with continuing use of the prosthesis.

The second remark concerns the definition of the ulcers. Ulcers are often seen in amputees using prosthesis (Meulenbelt et al. 2006, 2007). The definition of an ulcer used by the authors is not a worldwide used definition (Berke 2004). There is, however, no accepted definition. In this study primary and secondary problems in wound healing are compared to the development of an ulcer. Further on in this paper both problems are grouped together. An average surface area of the ulcer was measured to evaluate the possible effect of the intervention. It is uncertain this measurement of average surface should be used as an outcome measure of the success of an intervention because delayed wound healing and the development of ulcers are pooled together and some patients had several ulcers. As the authors already stated in their discussion as a weakness of their study, the depths of the ulcers were not measured, while this is an important measure in classification of ulcers (Wagner 1981). Finally the type of ulcer was not described.

The last remark is related to the categorization of limb used during the survey period. This subject is not elaborated in this paper. Only five patients were advised to discontinue prosthetic limb use; they continued to use their prosthesis which in two cases led to deterioration of the ulcer. What about the other three cases? In the meantime new patients received new prostheses or received socket adaptations.

With respect to Salawu et al.'s findings which touch on a very important issue in the rehabilitation of amputees, clarification to our remarks will help us better understand their results.

Henk E. Meulenbelt  
Jan H. B. Geertzen  
Center for Rehabilitation,  
University Medical Center Groningen and  
Northern Center for HealthCare Research,  
University of Groningen, The Netherlands

## References

- Berke G. 2004. Post operative management of the lower extremity amputee-standards of care. Official findings of the Consensus conference. *J Prosthet Orthot* 16(35):6–12.
- Meulenbelt HE, Dijkstra PU, Jonkman MF, Geertzen JHB. 2006. Skin problems in lower limb amputees: A systematic review. *Disabil Rehabil* 30:28(10):603–638.
- Meulenbelt HE, Geertzen JHB, Dijkstra PU, Jonkman MF. 2007. Skin problems in lower limb amputees: An overview by case reports. *J Eur Acad Dermatol Venereol* 21(2):147–155.
- Saluwa A, Middleton C, Gilbertson A, Kodavali K, Neumann V. 2006. Stump ulcers and continued prosthetic limb use. *Prosthet Orthot Int* 30:279–285.
- Wagner FW. 1981. The dysvascular foot: A system of diagnosis and treatment. *Foot Ankle* 2:64–122.